



Self-measured blood pressure technique:

How to take your own blood pressure

Before you measure

- 1. Use a certified, automated device to measure your blood pressure (BP) using your arm (not finger or wrist).
- 2. Use a cuff that is the right size for your arm.
- 3. Do not exercise, smoke, eat a large meal, take decongestants or have caffeine within 30 minutes of measuring your blood pressure.
- 4. Use the bathroom if you need to, before the measurement.
- 5. Rest for five minutes before measuring your blood pressure.

Position yourself correctly

- 6. 1 Sit in a chair, with your back supported.
 - 2 Sit with your legs uncrossed and feet flat on the floor (or stool).
 - Rest your arm on a table close to heart level.
 - 4 Place the blood pressure cuff over bare skin, on mid-arm at heart level and just above your elbow.

Perform blood pressure measurement

- 7. Do not talk, text, read, watch TV or use your phone, computer or tablet while measuring your blood pressure.
- 8. "Power on" the machine and push the start button.
 - When the machine stops, write down the upper and lower BP numbers (systolic and diastolic) if the machine does not store them automatically.
 - Wait one minute and then repeat (some machines will do this automatically). You should always check at least two blood pressure measurements one minute apart and write them down.
- 9. Take your two blood pressure measurements in the morning and two in the evening for one week, and report them to your doctor's office.





Self-measured blood pressure monitoring at home – flow sheet

Name:	Date of birth:
Name.	Date of birtin.

Instructions for self-measured blood pressure at home

Decide with your doctor or care team if you should use this form. You may not need to use it if your blood pressure device is able to store your readings and you are able to share those readings with your clinician.

(See "Self-measured blood pressure at home" handout for additional information.)

Rest for five minutes before measuring the first blood pressure

- 1. Take at least two measurements each time you check your blood pressure and write them down. Wait at least one minute between each measurement.
- 2. Write any factors you feel may have affected your blood pressure in the comments section.
- 3. Do this two times a day—once in the morning and once in the evening.
- 4. Give these numbers to your doctor or clinical office staff in person, during a telephone call or through secure computer messaging.

Morning			Evening			
Date	#1	#2	Comments/average	#1	#2	Comments/average
Day 1						
Day 2						
Day 3						
Day 4						
Day 5						
Day 6						
Day 7						

Morning			Evening			
Date	#1	#2	Comments/average	#1	#2	Comments/average
Day 1						
Day 2						
Day 3						
Day 4						
Day 5						
Day 6						
Day 7						

Copyright 2015 American Medical Association and The Johns Hopkins University. All Rights Reserved. Please feel free to adapt this content to meet the needs of your organization to improve health outcomes solely for non-commercial purposes. Use of the names of the American Medical Association or The Johns Hopkins University or the logos of either beyond use of the names in the required permission statement is not permitted on any adaptations without prior written authorization. The following should appear on the first page or screen of adaptions:

"This [content description, e.g., protocol or checklist] was adapted with permission of the American Medical Association and The Johns Hopkins University. All Rights Reserved. The original copyrighted content can be found at [link]."

